

A Guide

FOR **PARENTS AND CARERS** of children
aged **birth-5 years**


Common childhood illnesses

NHS
Islington
Clinical Commissioning Group

 ISLINGTON



This handbook has been produced by
Public Health for Camden and Islington
view online at www.islington.gov.uk/childillness





Welcome



This book has been put together by Public Health for Camden and Islington.

Every parent or carer wants to know what to do when a child is ill - use this handbook to learn how to care for your child at home, when to call a Doctor and when to contact Emergency Services.



Most of the problems you will come up against are simply an everyday part of growing up, often helped by a chat with your Midwife, Health Visitor or School Nurse. Almost all babies, toddlers and children will get the most common childhood illnesses like chickenpox, colds, sore throats and ear infections. While these are worrying at the time they are easily treated by your Doctor or at home with the support from a Doctor or Health Visitor rather than a trip to Accident & Emergency.



This handbook helps point you in the right direction and explains what you can do at home to help, or where you need to go to get assistance and advice. It has been put together with help from healthcare professionals. If you are worried you must get further advice. Trust your instincts, you know your child better than anybody else.

This handbook also contains general welfare information which will help you keep you and your child safe and healthy.

Visit www.islington.gov.uk/childillness to view this booklet online.



All factual content has been sourced from Department of Health, Birth to Five, 2009 edition, NHS Choices, British Association of Dermatologists, Meningitis Now. This information cannot replace specialist care. You need to get specialist help if you are worried, you know your baby best.



To view this booklet online scan this code with your smartphone

Contents

» A guide to services	4	» Earache	28
» Know the basics	6	» Fever	30
» Being sick	8	» Meningitis	32
» Crying	10	» Bumps & bruises	34
» Nappy rash	12	» Burns & scalds	36
» Rashes & dry skin	14	» Asthma	38
» Sleeping	16	» Chickenpox & measles	40
» Sticky eyes & conjunctivitis	18	» Urticaria or hives	42
» Wheezing & breathing difficulties	20	» Household accidents	44
» Upset tummy	22	» A healthy weight	46
» Constipation	24	» Worried about a child	48
» Coughs, colds & flu	26	» Good mental health	50
		» Useful contacts	52

A guide to services

We have a wide range of healthcare and children and family services.
See which service or professional is best to help you.

Self-care

Many illnesses can be treated in your home by using over the counter medicine from your Pharmacist and getting plenty of rest. Self-care is the best choice to treat very minor illnesses and injuries. If you are still worried contact NHS 111 or your Doctor.



111

If you think you need help urgently during the day or night you should call 111 before you go to any other health service. By calling 111 you will be directed straight away to the local service that can help you best. It is available 24 hours a day, 365 days a year and is free to call, including from a mobile. You should call NHS 111:

- When you need help fast but it's not life threatening.
- When you think you need to go to A&E or another NHS urgent care service.
- When it's outside of your Doctor's surgery hours.
- When you do not know who to call for medical help.
- If you do not have a local Doctor to call.

Minor Ailments Scheme

The Minor Ailments Scheme is available to all - it allows Pharmacists to manage simple common medical problems and issue medicines as if they had been prescribed by the GP. You do not always need to see your Doctor. Pick up a Minor Ailments Scheme form from your surgery reception. You will be issued with a Minor Ailments Passport. Take it to your local pharmacy who will be able to help you with things like:

- Colds and flu
- Constipation
- Earache
- Headlice
- Insect bites and stings
- Minor burns or lacerations
- Teething

Pharmacist

Your local Pharmacist will know about most everyday health issues. They can suggest the best medicine to help. There are often Pharmacists in supermarkets and many are open late. If your child has a temperature which has not come down with paracetamol or ibuprofen see your Doctor.

Doctor or GP

You will need to register with a local Doctor. Your Doctor can advise, give you the medicines you need and point you in the right direction if you need other specialist services. You will usually need to make an appointment. All Doctors will see a child quickly if you are worried. After 6.30pm weekdays, at weekends and public holidays you can call the GP out-of-hours service on NHS 111.

Health Visitor

The Health Visiting Team are there to support you when you need them. They will visit you at home or see you in a clinic. They can tell you where to get extra help if you need it. They are part of a team who are there to support you during the early years.

Children's Centres

Children's Centres are for families with children under five and many of the services are free. Together the centres offer a wide range of services including:

- Health Visitors
- Midwifery services
- Play sessions for children
- Parenting support (including support for teenage parents)
- Services for disabled children
- Speech and language support
- Healthy eating and weight management sessions.

Many centres also provide high quality early learning and childcare.

Urgent Care Centres

Urgent Care Centres treat patients who have an injury or illness that needs help quickly, but is not a 999 emergency. To book an appointment with a local Urgent Care Centre call 111.

A&E

For immediate, life-threatening emergencies, please call 999.

A&E and 999 are emergency services that should only be used when babies and children are badly injured or show symptoms of critical illness such as choking, chest pain, blacking out, or blood loss.

If a child in your care is ill or injured, choose from the following services available:

Grazed knee, Sore throat Coughs and colds, Mild tummy pain or headache	Self Care	You can treat minor illnesses and injuries at home by using the recommended medicines and making sure they get plenty of rest www.nhs.uk .
As a parent if you are: Unsure Confused Need help	 NHS 111 For 24 hour health advice and information.	Ring NHS 111 when it is less urgent than 999 Tel: 111 www.nhs.uk/111
Mild diarrhoea Mild skin irritations (including spots/rash) Mild fever	Pharmacist For advice on common illnesses, injuries and medication.	To find your local pharmacy and its contact details visit: www.nhs.uk/chemist (see Minor Ailments Scheme page 4).
High temperature Head injuries not involving loss of consciousness Persistent cough Worsening health conditions (inside GP hrs)	Doctor For the treatment of illnesses and injuries that will not go away.	Write your Doctor's (family doctor) telephone number here: <input type="text"/>
Minor bumps, cuts and possible fractures (during 9-5) Dehydrated Headache Tummy pain	Health Centre For treatment of minor illnesses and injuries without an appointment.	Health Centre or Walk-in Centre
Unexpected and sudden sickness Severe pain Worsening health conditions (outside GP hrs)	Urgent Care When you need healthcare in a hurry 24 hours a day.	A&E/Urgent Care Centre
Choking Loss of consciousness Fitting Broken bones	A&E or 999 For very severe or life threatening conditions.	A&E

NHS 111 is free to call from any landline or contract mobile phone. Pay-as-you-go mobile phones require 1 pence credit to make a call.

Know the basics

Being prepared and knowing the signs

Parents are usually good at noticing when something is wrong with their baby from quite early on. It is normal to worry that you won't recognise the signs that your baby is unwell. Trust your instincts, you know your baby best.

Learn how to spot the signs of serious illness and how to cope if an accident happens. If you know the basics and you are well prepared, you will find it easier to cope - and less scary. Keep a small supply of useful medicines in a locked cabinet or somewhere up high where a child cannot reach them. There is a useful list in the Pharmacist says box on the right, of things to have at home just in case. Make sure you've got the right strength of medicine for the age of your child, always follow instructions carefully and check use by dates. Read the label carefully. Children under 16 years should not take aspirin, unless on the advice of a Doctor. Aspirin use in children has been linked with Reye's syndrome. This is a very rare but fatal condition that affects the brain and liver.

If your baby seems to have a serious illness it is important to get medical attention as soon as possible.

1

My baby is crying more than usual and seems to be irritable and hot.

2

Is there a fever, have you checked their temperature? Have you tried paracetamol? Remember to check the label to give the right dose.

3

If you have tried this and it has not worked see your Pharmacist. **If temperature is 38°C or above coupled with a rash**, contact your Doctor immediately.

Source: NHS choices



Pharmacist says

Keep a small supply of useful medicines. Include things like:



Thermometer



Plasters



Liquid painkillers (e.g. paracetamol or ibuprofen). Read label carefully.



Barrier cream



Natural oils like vegetable oil (for dry skin)



Health Visitor says

Possetting is 'normal' during or after a feed. If this carries on at other times, between feeds it may be a tummy bug. It is important for babies to have plenty of fluids to stop any dehydration.

Being sick

A problem likely to get better on its own

It is common for babies to be sick, often in the early weeks as they get used to feeding and their bodies develop. Possetting is when your baby brings up small amounts of milk; when your baby vomits this will be a much larger amount. It can be frightening for your baby, so they are likely to cry. Lots of things can cause your baby to be sick.

Make sure your baby is positioned correctly when breast or bottle feeding as incorrect positioning can cause a baby to be sick.

Being sick often or lots of it, may be due to 'gastric reflux' where acid from the tummy can come up again. Babies can be grumpy and it can sometimes lead to poor feeding. If your baby is feeding well but doesn't seem themselves, you may just need to change the baby's position during a feed to make them more upright. Feeding smaller amounts and more often may also help.

1

I have a newborn. I have just given my baby a feed.

2

They always seem to bring up small amounts of milk.

3

This is known as 'possetting'. As they develop it will stop naturally. Talk to your Health Visitor.

Doctor says

After the first few months, if your baby is suddenly sick it is more likely to be caused by a stomach virus rather than possetting. Gastroenteritis is a tummy bug (see [Upset tummy page 22](#)), which can come with diarrhoea (runny poo).

This is more serious in babies than older children because babies easily lose too much fluid from their bodies and become dehydrated. If they become dehydrated they may not pass enough urine, lose their appetite and have cold hands and feet.

Get expert advice. If your baby is unwell, or if vomiting has lasted more than a day, get your Doctor's advice straightaway.



Health Visitor says

Know your baby. Try to understand what it is they need. Finding out why your baby is crying is often a matter of going through all the possible options.

Things to check first are:

- ✓ Does their nappy need changing?
- ✓ Could they be hungry?
- ✓ Could they be too hot?
- ✓ Could they be too cold?
- ✓ Does their cry sound different?

These are simple things which could be causing your baby to cry.

Crying

Understanding why

All babies cry, especially in the first few weeks after birth. Crying is their way of letting you know they need something or are uncomfortable. They may need changing, they may be hungry or just need a cuddle. Always burp your baby after a feed as this will help.

Look out for signs that your baby is trying to tell you they are hungry. Early signs are things like putting their hands to their mouth, becoming restless and stretching. By recognising these cues you may avoid hunger crying altogether and the need to calm baby down before a feed.

If your baby cries suddenly and often, but they otherwise appear to be happy and healthy, they may have colic. Colic is common and although uncomfortable it is not serious and usually affects babies only in the first few months of their lives. The most common symptom of colic is continuous crying, which typically occurs in the late afternoon or evening. Other signs include a flushed appearance, drawing their legs to their chest, clenching fists, passing wind and trouble sleeping.

When a baby cries, it can be upsetting. It is very important to stay calm and don't be afraid to ask for help. Do not shake your baby.

1

My baby is crying more than usual.

2

Have you followed the advice given by your Health Visitor? Have you thought about what your baby is trying to tell you?

3

If you have tried this and it has not worked speak to your Health Visitor, or contact your Doctor if you are worried.



Doctor says

If your baby's crying seems different in any way (such as a very high-pitched cry or a whimper), then seek medical advice. Crying can sometimes be a sign that your baby is unwell. Trust your instincts - you know your baby best.



Health Visitor's cradle cap tips

This is the name given to the yellowish, greasy scaly patches on the scalp of newborns and usually appears in the first 3 months. It can look like a bad case of dandruff and clears up over time without causing your baby discomfort.



Wash scalp gently every day using luke warm water.



Use a small amount of natural oil (vegetable oil) on the scalp and leave on for 15 minutes before washing off with luke warm water.

It is important not to pick at the scales as this may cause infection.

Health Visitor's nappy rash tips



Leave your baby in a warm, safe place with no clothes or a nappy on, to let the air get to their skin.



Use a barrier cream. (see Pharmacist says box opposite).



Remember to change and check their nappy often.

Nappy rash

A common problem that's easy to treat

Nappy rash is very common and can affect lots of babies. It is usually caused when your baby's skin comes into contact with wetness that collects in their nappy.

A nappy rash causes your baby's skin to become sore. The skin in this area may be covered in red spots or blotches. You might need to change their nappy more often.

Most nappy rashes can be treated with a simple skincare routine and by using a cream you can get from the Pharmacist. With a mild nappy rash, your baby won't normally feel too much discomfort.

Dry skin

A baby's skin is thinner and needs extra care. Dry, flaky skin, some blemishes, blotches and slight rashes are normal in newborns and will naturally clear up. If your baby is otherwise well but has a rash you are worried about contact your Midwife or Health Visitor.

1

There is a red, sore rash around the nappy area. Baby is uncomfortable and cries a lot.

2

Has baby been in a dirty nappy for a long time? Have you followed advice from your Health Visitor, or spoken to your Pharmacist?

3

Change nappy often. Speak to your Health Visitor and if you are worried see your Doctor.



Pharmacist says

Call in and talk to us about creams we can provide you with over the counter.

There are two types of nappy cream available. One is a barrier cream to keep wetness away from your baby's skin. The other is a medicated cream, that is good for clearing up any soreness but should only be used when advised by a health professional.



Health Visitor says
Baby skin is more delicate than ours. Try to limit the amount of products you use on their skin and never leave your baby out in the sun.

Rashes & dry skin

Baby skin needs extra care

A baby's skin is thinner and needs extra care. Dry, flaky skin, some blemishes, blotches and slight rashes are normal in newborns and will naturally clear up. If your baby is otherwise well but has a rash you are worried about contact your Midwife or Health Visitor. Another common rash for babies is heat rash. This mainly appears on the head and neck as tiny red spots and is nothing to worry about. Keep them warm but not hot and try to dress them in natural cotton clothes, with nothing that can rub on their skin.

Your baby may also suffer from something called cradle cap. This is the name given to the yellowish, greasy scaly patches on the scalp of newborns and usually appears in the baby's first three months. It can look like a bad case of dandruff and is harmless, it doesn't cause any irritation to your baby and usually clears up by the time they are two years old.

1

Your baby's skin may be flaky and dry.

2

Dry skin is common in newborn babies, as their skin is 15 times thinner than that of an adult.

3

Avoid soap and using products on your baby's skin. Wash your baby in clean, warm (not hot) water.



Doctor says
Go to A&E immediately if your baby has a rash that does not disappear when you press a glass to it. This may be a sign of meningitis and needs to be seen by a Doctor no matter how well your baby seems.
Seek immediate advice if your baby has a rash and a high temperature or vomiting.

The above information cannot replace specialist treatment. If you are worried call NHS 111 or contact your Doctor.



Safer sleeping

Keeping your baby safe and healthy:

- Place your newborn baby on their back to sleep, in a cot in your bedroom for the first six months.
- Place your newborn baby in the 'feet to foot' position i.e. baby's **feet** at the **foot** of the bed/cot. Once babies move around and roll, they will find a position that's comfortable for them.
- Do not let your baby get too hot and keep his head uncovered.
- Never sleep with your baby on a sofa or armchair.
- Do not smoke in pregnancy or let anyone smoke in the same room as your baby.

It's dangerous for your baby to sleep in your bed if you (or your partner):

- Are a smoker (even if you never smoke in bed or at home).
- Have been drinking alcohol or taken any drugs.
- Have taken any medication that makes you drowsy.
- If your baby was premature (born before 37 weeks).
- If your baby was low birth weight (less than 2.5kg or 5 1/2 lb).
- If you or your partner are overweight.

It is very dangerous to fall asleep together on a sofa, armchair or settee and it is also risky to allow a baby to sleep alone in an adult bed.



Sleeping

Patience, praise and peace

There are many different reasons why babies and toddlers do not sleep through the night. Feel confident in yourself to know whether your child is really distressed or just restless. During the first months babies are often wakeful at night and sleep more lightly.

Try to establish a regular day and night time sleep routine for your child until they are 2-3 years old. Put them to bed at a regular time, routine is important. Prepare a warm, comfortable place for them to relax in. Reading to your child at bedtime helps them to unwind, and gives you some special time together. If your child is scared of the dark, try keeping a night light on. Lack of sleep can cause a child to become hyperactive and have extremes in behaviour. Only breastfeeding babies should ever be fed in an adult bed, and if so, should be positioned on the outside of the bed and returned to the cot after the feed has finished.

Bedwetting may be stressful for both of you and can wake your child. It is not easy to know why some children take longer to be dry at night than others. Try not to lose your patience or punish them, your child is not doing this on purpose. Children learn at their own pace and praise and support will help.

1

I am so tired when my baby wakes up at night it seems easier to share the bed.

2

The safest place for your baby to sleep is in a cot by your bedside for at least the first 6 months. Try to establish a regular sleep routine.

3

Speak to your Health Visitor about how to keep baby safe and get some sleep.



Health Visitor

If your sleep is often disturbed, arrange for a trusted relative or friend to care for your baby or child so that you can get some sleep. Talk to your Health Visitor. Your child's sleep may be disturbed by bedwetting. Between the ages of three and four they are likely to have the occasional accident but gradually more and more nights will be dry.



Sticky eyes & conjunctivitis

Two different issues

'Sticky eyes' are common in newborn babies and young children while their tear ducts are developing. You may see some sticky stuff in the corner of the eyes or their eyelashes may be stuck together.

It normally clears up on its own, but you may have to clean your baby's eyes regularly with damp cotton wool. Use clean, cooled boiled water or breast milk.



Wipe each eye from the corner by the nose outwards. Use a clean piece of cotton wool for each wipe. Remember to wash your hands before and afterwards and avoid sharing towels to prevent spreading infection. Gentle massaging round the tear duct might help to open the tear duct to allow discharge to drain away.

1

Is there discharge in the corner of your baby's eye and do their eyelashes appear to be stuck together?

2

Sticky eyes is a common condition that affects most babies, speak to your Health Visitor.

3

Use cooled boiled water on a clean piece of cotton wool for each wipe.



Doctor says - Conjunctivitis

The signs of 'sticky eyes' can sometimes be confused with an infection called 'conjunctivitis'. With conjunctivitis the signs are yellowy, green sticky goo which comes back regularly. If you notice this ask your Pharmacist for advice and treatment. Conjunctivitis can be passed on easily, so wash your hands and use a separate towel for your baby.

Source: DoH 2006.

Bronchiolitis

Bronchiolitis is a common respiratory tract infection that affects babies and young children under a year old. The early symptoms are similar to those of a common cold and include a runny nose and cough.

As it develops, the symptoms of bronchiolitis can include: A slight fever, a persistent cough and difficulty feeding.

Symptoms usually improve after three days and in most cases the illness isn't serious. However, contact your Doctor or Health Visitor if your child is only able to feed half the normal amount or is struggling to breathe, or if you are generally worried about them.

Source: www.nhs.uk/conditions/Bronchiolitis/



1

My child with croup has a distinctive barking cough and makes a harsh sound, when they breathe in.

2

Comforting your child is important as symptoms may worsen if they are agitated or crying. Mild cases of croup can be managed at home. If your child has a fever, children's paracetamol will help lower their temperature.

3

Some people have found that allowing their child to breathe in steam from a hot bath or shower in a closed room has eased symptoms. Take care as there is a risk of scalding if your child is left alone. If symptoms get worse contact your Doctor.

Wheezing & breathing difficulties

Look at the signs

Any kind of breathing difficulty your infant or child experiences can be scary for parents. It may be nothing to worry about and could just be normal baby 'snuffles'.

Use your instincts with newborns and babies. It could be:

- Rapid breathing or panting, which is common. There is no other sign of illness, it comes and goes and your baby is breathing comfortably most of the time, there's normally no need to worry.
- Breathing may sound a bit rattly. Try holding your baby upright.
- Occasional, coughing or choking which may occur when a baby takes in milk too quickly with feeds. Try to slow things down a bit. Check feeding position.
- A cold or mild cough. Keep an eye on them at this stage and use your instincts. If you are worried talk to your Health Visitor.

In older babies and toddlers you may notice:

- Coughing, runny nose, mild temperature - (see page 26 Coughs, colds & flu).
- Croup (hoarse voice, barking cough) needs to be assessed by a Doctor and may need treating with steroids.
- Child appears pale.

If you're worried about your child wheezing or having breathing difficulties even after reading this, contact your Doctor or call 999 immediately.

Source: NHS Choices - Symptoms of bronchiolitis



Doctor's tips

Get help and contact your Doctor now if your child:

- ✓ Seems to find breathing hard work and they are sucking in their ribs and tummy.
- ✓ They can't complete a full sentence without stopping to take a breath.

Get help and call 999 or take them to A&E now if:

- ✓ Their chest looks like it is 'caving in.'
- ✓ They appear pale or even slightly blue-ish.



Pharmacist says

There are lots of ways you can care for your child at home. Things to try are:

- ✓ Give them regular drinks - try small amounts of cold water. Breastfeed on demand if breastfeeding.
- ✓ Being extra careful with hand hygiene (use soap and water or anti-bacterial hand gel and dry hands well with a clean towel).
- ✓ Rehydrating solutions come in pre-measured sachets to mix with water. It helps with dehydration.

If your child is unwell for more than 24 hours speak to your Doctor. If your baby is newborn or very unwell contact your Doctor straight away.

Upset tummy

Not nice for you or your baby

Sickness and diarrhoea bugs are caught easily and are often passed on in places where there are lots of children.

Feeling sick and suddenly being sick are normally the first signs. Diarrhoea can follow afterwards. If your child is not vomiting frequently, is reasonably comfortable in between and you are able to give them frequent small amounts of water, they are less likely to become dehydrated and probably don't need to see a Doctor.

Speak to your Doctor if they are unwell for longer than 24 hours or sooner if they are newborn or if you notice signs of dehydration. →

If you're breastfeeding, keep on doing so even more frequently. Offer older children plenty of water, or an ice-lolly for them to suck. If they want to eat, give them plain foods like pasta or boiled rice (nothing too rich or salty).

Keep them away from others, especially children, who may pick up infection. Be extra careful with everyone's handwashing.

1

My baby has diarrhoea and is being sick.

2

Have you given them lots of water? This will help prevent them becoming dehydrated if it is a tummy bug. Speak to your Pharmacist and ask about a rehydrating solution.

3

Speak to a Doctor if symptoms show no sign of improvement after 24 hours or straight away if they are newborn.

Signs of dehydration

- ✓ Sunken fontanelle (i.e. the soft spot is more dipped in than usual).
- ✓ Less wet nappies (i.e. they wee less).
- ✓ More sleepy than usual.
- ✓ Dry mouth.

Try rehydrating solution from your Pharmacist.

Breastfed babies

In the first 4-6 weeks, breastfed babies should have at least 2 soft, yellow poos per day. After this time they may need to go less often, but the poo should still be soft.



Constipation

Common and easy to treat

Constipation means passing hard poos, with difficulty and less often than normal. Babies can have a poo several times a day or once every few days. Babies often appear to strain and go red in the face when having a poo - this is normal. In children, having a poo anything from 3 times a day to once every other day can be normal so long as the poos are soft and easily passed. Sometimes bad constipation can cause a child to dirty their pants.

Doctors do not understand what causes most constipation in children but it is important to get it treated because if a child is constipated they may find it painful to go to the toilet. This creates a vicious circle; the more it hurts, the more they hold back, the more constipated they get. Once a child is really constipated they may stop wanting to go to the toilet altogether. Take your Minor Ailments Passport to the Pharmacist for advice and treatment (see page 5). If you are still worried, talk to your Doctor.

1

Has your child stopped wanting to go to the toilet altogether?

2

If your child is constipated, they may find it painful to go to the toilet.

3

Ask your Health Visitor or Pharmacist whether a suitable laxative may help.

Source: NICE guidelines 2009, Constipation in children

Health Visitor says

To avoid constipation and help stop it coming back make sure your child has a balanced diet including plenty of fibre such as fruit, vegetables, baked beans and wholegrain breakfast cereals. We do not recommend unprocessed bran (an ingredient in some foods), which can cause bloating, flatulence (wind) and reduce the absorption of micronutrients. Drink plenty of fluids.

If a bottle fed baby becomes constipated you can try offering water between feeds (never dilute baby milk). Check that you are making up the powdered feed correctly and that you are not overfeeding. First Stage Formula is best until a year when they can go on to cow's milk. Formula for 'hungrier babies' can lead to constipation. If the problem doesn't go away, talk to your Health Visitor or Doctor again.

1

My child keeps coughing and sneezing, has a mild temperature and seems generally unwell.

2

Have they recently started nursery? Catching colds is very common. Have you spoken to your Pharmacist about junior paracetamol and cough medicines?

3

If symptoms last for more than 10 days or your child is coughing up yellow 'goo' they may have an infection. Contact your Doctor.

Paracetamol - can be given to children for pain or fever. Check you have the right product, dose and strength for your child's age. Read the box carefully.

Ibuprofen - can be given to babies and children of 3 months and over who weigh more than 5kg. Read the box carefully. Avoid if your child has asthma unless advised by your Doctor.

Do not give paracetamol and ibuprofen at the same time.

If your child is still distressed you could consider changing to the other drug when the next dose is due.

Aspirin - do not use for children under 16.

Don't pass it on:

Catch it Germs spread easily. Always carry tissues and use them to catch coughs or sneezes.

Bin it Germs can live for several hours on tissues. Dispose of your tissue as soon as possible.

Kill it Hands can pass on germs to everything you touch. Clean your hands as soon as you can.

Coughs, colds & flu

Not usually serious

You will probably find when your child goes to playgroup or nursery that they get lots of coughs, colds and sniffles. There are some good things about this though as it helps the body build up a natural immune system.

Flu can be more serious than a cold and leave your child feeling quite unwell. Flu tends to come on more suddenly and severely than a cold. Your child may feel achy and uncomfortable, and be ill for a week or more.

Most bugs will run their course without doing any real harm because they will get better on their own. An annual nasal spray flu vaccine is available as part of the NHS Childhood Vaccination Programme. Ask your Health Visitor for details.

Things you can do at home to help:

- ✓ Give your child lots to drink.
- ✓ Try paracetamol (not aspirin).
- ✓ Keep them away from smoke and anyone who smokes.
- ✓ Talk to your Pharmacist but remember that coughing is the body's way of keeping the lungs clear. Use the Minor Ailments Passport (see page 5).

See your Doctor if:

- ✓ Your baby has a temperature of 38°C or more.
- ✓ They have a fever with a rash.
- ✓ They are not waking up or interacting.
- ✓ Your child is finding it hard to breathe.

Source: 2013 NICE guidance.



Pharmacist says

Children can often be treated using over the counter medicines to help to bring down a raised temperature. Use the Minor Ailments Passport (see page 5). Paracetamol can help. Check the label carefully. Some are available as a liquid for children and can be given from the age of about three months. Check with the Pharmacist and tell them how old your child is. Flu symptoms are more severe and you may need to see your Doctor.



Earache

A baby's ears need to be treated with care

Ear infections, which can result in earache are common in babies and toddlers. They often follow a cold and can sometimes cause a temperature. A child may pull at their ear, but babies often cannot tell where their pain is coming from, so they just cry and seem generally uncomfortable.

Babies have some natural protection against infections in the first few weeks - this is boosted by breastfeeding. In babies and toddlers, bacteria pass from the nose to the ears more easily. Ear infections can be painful and your child may just need extra cuddles and painkillers. Take your Minor Ailments Passport to the Pharmacist for advice and treatment (see page 5). Your child may have swollen glands in their neck - this is the body's way of fighting infection.

Children who live in households where people smoke (passive smoking) or who have a lot of contact with other children, like those who go to nursery, are more likely to get ear infections. Bottle fed infants are more prone to ear infections, due to the sucking technique. Try to cut down/discontinue bottles and discuss with your Health Visitor.

1

My toddler has earache but seems otherwise well.

2

Have you tried infant paracetamol or ibuprofen from your Pharmacist? Do not put oil or cotton buds into your child's ears.

3

Most ear infections get better by themselves. Speak to a Doctor if symptoms show no sign of improvement after 24 hours, your child seems in a lot of pain or you notice fluid coming from the ear.

Source: DoH Birth to five edition 2009.



Health Visitor's tips

- ✓ A baby's ears need to be treated with care when cleaning.
- ✓ Never use a cotton bud inside your child's ear.
- ✓ If they have a temperature wax may ooze out.
- ✓ Use different, clean damp cotton wool on each ear to gently clean around the outer area.

What are the signs of an ear infection?

The signs are a raised temperature, general irritability and pain or discomfort. The ears may be red and your baby may pull them because they are uncomfortable. They may even have a pus-like discharge, which can also be associated with a blocked feeling in the ear or hearing loss. Although most ear infections settle down without any serious effects, there can be mild hearing loss for a short time (two to three weeks).



Babies under 6 months:

- A baby younger than 3 months with a temperature of 38°C or more without a rash needs to be assessed by a Doctor.
- A baby older than 3 months with a temperature of 39°C or more without a rash needs to be assessed by a Doctor.
- A baby of any age with a temperature of 38°C and a rash should be seen immediately as stated.

Fever

Common in young children

If your child has a fever, he or she will have a body temperature above 38°C. Your child may also feel tired, look pale, have a poor appetite, be irritable, have a headache or other aches and pains and feel generally unwell.

A fever is part of the body's natural response to infection and can often be left to run its course provided your child is drinking enough and is otherwise well. It is important to prevent your child from becoming dehydrated, which can lead to more serious problems. As a guide, your child's urine should be pale yellow - if it is darker, your child may need to drink more fluids.

Fevers are common in young children. They are usually caused by viral infections and clear up without treatment. Take your Minor Ailments Passport to the Pharmacist for advice and treatment. However, a fever can occasionally be a sign of a more serious illness such as a severe bacterial infection of the blood (septicaemia), urinary tract infection, pneumonia or meningitis.

Always seek medical advice if your child develops a fever soon after an operation, or soon after travelling abroad.

1

My toddler is hot and grumpy and seems upset.

2

Have you tried liquid paracetamol or ibuprofen? Do not give both at the same time. Have you made sure they are drinking lots of fluids?

3

If their temperature remains over 38°C and doesn't come down, contact your Doctor.



Older children:

A little fever isn't usually a worry. Contact your Doctor if your child seems unusually ill, or has a high temperature which doesn't come down.

- It is important to ensure your child drinks as much fluid as possible.
- Keep the room at a comfortable temperature (18°C).
- Give liquid paracetamol or ibuprofen in the correct recommended dose for your child. Do not give both at the same time.
- Aspirin should not be given to children under 16.



The glass test

The glass test is a really useful way of spotting suspected meningitis. If your child has a cluster of red or purple spots, press the side of a clear drinking glass firmly against the rash.

Go straight to the Accident and Emergency Department



In this example the spots are still visible through the glass. Contact a Doctor immediately. If you cannot get help straight away go to A&E.



In this example the spots under the glass have virtually disappeared. It is unlikely to be meningitis but if you are still worried contact NHS 111, your Doctor or go to A&E.

Source: Meningitis Now

Meningitis

A serious, contagious illness

Babies and toddlers are most vulnerable as they cannot easily fight infection because their immune system is not yet fully developed. They can't tell you how they are feeling and can get a lot worse very quickly. Keep checking them.

Meningitis is a swelling around the brain. It's a very serious, contagious illness which can sometimes get confused with other more common illnesses, but if it's treated early most children make a full recovery.

You should always treat any case of suspected meningitis as an emergency.

Early signs may be like having a cold or flu. Children with meningitis can become seriously ill very fast, so make sure you can spot the signs. Your child may have a cluster of red or purple spots. Do the glass test. This rash can be harder to see on darker skin, so check for spots over your baby or child's whole body as it can start anywhere (check lightest areas first). **However, the rash is not always present - be aware of all the signs/symptoms.**

The presence of fever and any other of the above symptoms should be taken extremely seriously. Not all children will show all the signs listed on the right.

1

My child is showing some of the signs of meningitis.

2

Have you tried the glass test?

3

Treat all cases of suspected meningitis as an emergency. If the spots do not fade under pressure **call 999 or go to A&E.**

If you are worried, contact a Doctor. If you cannot get help straight away go to A&E.



Doctor says

If any of the signs below are present contact a Doctor.



Fever, cold hands and feet



Floppy and unresponsive



Drowsy and difficult to wake



Spots/rash. Do the glass test



Rapid breathing or grunting




Fretful, dislikes being handled



Unusual cry or moaning





Keeping them safe

Being a toddler means your child is discovering the world around them. This can result in bumps and bruises. It is almost impossible to prevent every accident although there are things we can do at home which might help.

Bumps & bruises

Part of growing up

Minor cuts, bumps and bruises are a normal part of growing up. Allowing your child to explore the world around them (with supervision) helps them develop and learn. Most of your toddler's bumps will require no more than a cuddle to make them better. You will quickly be able to tell by the noise of the bang, the reaction of your child and the colour of the area affected, which are the more serious bumps. If your child has unexplained bruising or injury you need to find out how this happened.

If it looks like the bump may swell then use a cold flannel (soaking the cloth with cold water) or ice pack (but don't put ice directly onto the skin) to help reduce swelling and to cool the area for at least a few minutes.

If your child has had a bump to the head and it looks serious or symptoms worsen call a Doctor. Read the information on the right. ➡

If your child is under a year old and has a bump on the head get advice from a Doctor.

1

After a fall comfort the child, check for injuries, treat bumps and bruises.

2

Give the child some painkillers and let the child rest whilst watching your child closely.

3

Seek immediate help if:

- They have seriously injured themselves.
- They are unconscious.
- They have difficulty breathing.
- They are having a seizure.

Head injury

One of the signs of a severe head injury is being unusually sleepy, this does not mean you cannot let your child sleep.


You need to get medical attention if:

- **They are vomiting persistently (more than 3 times).**
- **They are complaining it hurts.**
- **They are not responding at all.**
- **Pain is not relieved by paracetamol or ibuprofen.**

If they are tired from what's happened, or from crying, then it is fine to let them sleep. If you are worried in any way about their drowsiness, then you should wake your child an hour after they go to sleep.

Check that they are okay, and that they are responding normally throughout the night.

If you are still worried, contact NHS 111. If you cannot get help go straight away to the Accident and Emergency Department.



Burns & scalds

Knowing what to do

A burn is damage to the skin, which is caused by direct contact with something hot. Burns can also be caused by certain chemicals, electricity and friction. A scald is a burn that is caused by a hot liquid or steam. Scalds are treated in the same way as burns.

Treat any burn or scald straight after the accident but always take your child to hospital for anything more than a very small burn or scald. A baby's skin is very delicate and can be scarred without the right treatment.

Cool the burnt area by placing under cool running water for at least twenty minutes. When the burn has cooled, cover it with a sterile dressing, food quality cling film or a plastic bag. Don't wrap it too tightly. Give paracetamol or ibuprofen. Then take your child to hospital.

Remember to keep hot drinks out of children's reach.

1

Treat the burn or scald straight after the accident by running under cold water for 20 minutes.

2

Do not use creams, lotions or ointments on the burn or scald.

3

Always take your child or baby to hospital if it is anything other than a very mild burn.



Do

Hold the affected area under cold water for at least 20 minutes (make sure your child does not get too cold). Cover the burn with cling film, then wrap in a cloth soaked in cool water.

Don't

Apply fatty substances like butter or ointment as this will not help and will only waste time for hospital staff who will have to clean the area before it can be treated.

If you are still worried, contact NHS 111. If you cannot get help straight away go to the Accident and Emergency Department.



Health Visitor says

Smoking during pregnancy or around your child can increase risk of asthma. Breastfeeding for as long as possible can help reduce risk of getting asthma.

Symptoms of severe asthma

Symptoms include repeated coughing and wheezing, shortness of breath and bringing up phlegm. Symptoms often get worse at night.

Asthma

Know the symptoms

Asthma is a common long-term condition that can be well controlled in most children. The severity of asthma symptoms varies between children, from very mild to more severe. Parents learn how to be prepared and how to recognise symptoms and deal with them.

Asthma affects the airways and makes it difficult to breathe and causes wheezing, coughing, shortness of breath and can make the chest feel tight.

A sudden, severe onset of symptoms is known as an asthma attack. Asthma attacks can sometimes be managed at home but may require hospital treatment. They are occasionally life threatening.

Triggers can include exercise (especially in cold weather), an allergy with dust mites, animal fur, grass and tree pollen or exposure to air pollution, especially tobacco smoke or a cold virus. Asthma often runs in families. If your child is overweight, there is a greater likelihood that they could develop asthma. Talk to your Health Visitor about support from weight management services in the borough.

Call 999 to seek immediate medical assistance if your child has severe symptoms of asthma.

1

My child seems to wheeze and cough a lot, it seems to get worse at night.

2

Have you tried reducing any possible amounts of dust around the home? Do you smoke? Have you discussed with your Health Visitor?

3

If symptoms persist see your Doctor. If your child has a serious asthma attack **call 999**.



Doctor says

Your Doctor will normally be able to diagnose asthma by asking about your child's symptoms, examining their chest and listening to their breathing. They will want to know about your child's medical history and whether there is a history of allergic conditions in your family. They will also want to know about the circumstances surrounding the onset of your child's symptoms, such as when and where it happened, because this could help to identify the possible trigger(s) of their asthma.



Midwife says

If you are pregnant and have had chickenpox in the past it is likely that you are immune to chickenpox. However, please contact your Doctor or Midwife for advice.

Chickenpox & measles

Take rashes seriously

Chickenpox

Chickenpox is a mild disease that most children catch at some point. The spots often look like mosquito bites and can appear on any part of the body. After having chickenpox, the virus stays in the body. Later in life the virus can come back in a different form known as shingles.

Chickenpox is easy to pass on to someone who has not had it before. If your child has chickenpox keep them away from others.

Chickenpox can be incredibly itchy, but it's important for children to not scratch the spots so as to avoid future scarring. One way of stopping scratching is to keep fingernails clean and short. You can also put socks over your child's hands at night to stop them scratching the rash as they sleep.

If your child's skin is very itchy or sore, try using calamine lotion or cooling gels. These are available in pharmacies and are very safe to use. They have a soothing, cooling effect.

Painkillers

If your child is in pain or has a high temperature (fever), you can give them a mild painkiller, such as paracetamol (available over-the-counter in pharmacies). Always read the manufacturer's dosage instructions. Do not give aspirin to children under the age of 16.

Measles

Measles is a very infectious illness. About one in five children with measles experiences complications such as ear infections, diarrhoea and vomiting, pneumonia, meningitis and eye disorders. One in 10 children with measles ends up in hospital. There is no treatment for measles. Vaccination is the only way of preventing it. If your children haven't yet had the MMR vaccination, don't delay. Speak to your Health Visitor.

Once the rash starts, your child will need to rest and you can treat the symptoms until your child's immune system fights off the virus. If there are no complications due to measles, the symptoms will usually disappear within 7-10 days.

Closing curtains or dimming lights can help reduce light sensitivity.

Damp cotton wool can be used to clean away any crustiness around the eyes. Use one piece of cotton wool per wipe for each eye. Gently clean the eye from inner to outer lid.



Foods to avoid:

There is controversy over the role of diet in people with long-term hives. There are two groups of chemicals in some foods that may trigger urticaria. It is important to discuss your child's diet with your Health Visitor.

Avoid:

- Shellfish
- Strawberries, bananas, mangoes, pumpkin, tangerines, kiwi
- Tomatoes, peas
- Fish
- Chocolate
- Pineapple

Cut down on:

- Spices
- Orange juice
- Raspberries
- Tea

Source: Allergy UK



Urticaria or hives

Easy to confuse with more serious illnesses

Childhood rashes are very common and often nothing to worry about. Most rashes are harmless and go away on their own.

Urticaria or hives is a raised, red, itchy rash that appears on the skin. It can be frightening especially if you don't know the cause. It happens when a trigger causes a protein called histamine to be released in the skin. Histamine causes redness, swelling and itching, the rash can be limited to one part of the body or spread across large areas of the body. It can sometimes be confused with other types of more serious rashes such as meningitis.

Hives can be triggered by many things, including allergens (such as food or latex), irritants (such as nettles), medicines or physical factors, such as exercise or heat. But usually no cause can be identified. It's a common skin reaction that's likely to affect children. The rash is usually short lived and mild, and in many cases does not need treatment as the rash often gets better within a few days. If you're struggling with it, a medication called antihistamine usually helps. Creams help with the itching and are available over-the-counter at pharmacies. Speak to your Pharmacist for advice.

1

My child has developed itchy red spots.

2

It can be difficult to identify what has triggered the rash. Try to think about any new or different foods they have had.

3

If itching persists ask your Doctor about anti-histamine medication.



Doctor says

If your child has urticaria avoid:

- Food such as peanuts, shellfish, eggs and cheese.
- Environmental factors such as pollen, dust mites or chemicals.
- Insect bites and stings.
- Emotional stress.
- Some medications - do not stop any prescribed medicines before you speak to your Health Visitor or Doctor.
- Physical triggers such as pressure to the skin, change in temperature, sunlight, exercise or water.

Source: www.nhs.uk/conditions/skin-rash-children

Cuts

Glass causes serious cuts with many children ending up in A&E.

PREVENTION:

Do not leave drinking glasses on the floor. Make sure glass bottles are up high.

WHAT TO DO:

If the cut is not serious bathe the area, make sure there is no glass left and cover with a clean non-fluffy cloth.

If the cut is serious, is bleeding a lot or has a piece of glass under the skin (maybe they trod on some glass) go to A&E.

Drowning

Many children drown, often in very shallow water. It happens in the bath, in garden ponds, paddling pools and water butts.

PREVENTION:

Supervise children near water at all times. Use a grille on ponds and fill in a garden pond to use as a sand pit. Learn to swim.

WHAT TO DO:

Get your child out of the water. Try to get them to cough up any water. If they are not responding **call 999**.

Poisoning

Poisoning from medicines, household products and cosmetics are common.

PREVENTION:

Lock all chemicals, medicines and cleaning products away.

WHAT TO DO:

Find out what your child has swallowed and take it with you to A&E.

Strangulation

Window blind cords and chains can pose a risk for babies and children who could injure or even strangle themselves on the hanging looped cords.

PREVENTION:

- Install blinds that do not have a cord, particularly in a child's bedroom.
- Do not place a child's cot, bed, playpen or highchair near a window.
- Pull cords on curtains and blinds should be kept short and kept out of reach.
- Tie up the cords or use one of the many cleats, cord tidies, clips or ties that are available.
- Do not hang toys or objects that could be a hazard on the cot or bed.
- Don't hang drawstring bags where a small child could get their head through the loop of the drawstring.

WHAT TO DO:

Untangle child, contact the emergency services and start CPR.

Source: RoSPA

Household accidents

Keeping your child safe in and around the home

Falls

For babies the biggest danger is rolling off the edge of a bed, or changing surface. For toddlers it is more about falling from furniture or down stairs.

PREVENTION:

Make sure baby cannot roll off any surfaces, put pillows around them. Do not put a bouncing cradle or car seat on a surface where they could wriggle off.

Use stair gates for toddlers. Make sure balconies are locked and fit restrictors and safety locks to windows.

WHAT TO DO:

If your child has a serious fall dial 999.

Choking

Babies and toddlers can easily swallow, inhale or choke on small items like balloons, peanuts, buttons or plastic toy pieces or strings or cords.

PREVENTION:

Check on the floor and under furniture for small items. Find out more about CPR (a first aid technique that is a combination of rescue breaths and chest compressions. Sometimes called the 'kiss of life').

WHAT TO DO:

If your child is choking act immediately and calmly. Make sure you do not push the object further down the throat. Encourage your child to cough. Use back blows, if they become unconscious call for help (do not leave your child alone), start CPR, **call 999**.
www.redcrossfirstaidtraining.co.uk

Burns, scalds and fires

Domestic fires are a risk to all children. Never let a child play with lighters, matches or candles. Burns from hair straighteners and household appliances are common. Hot drinks cause burns and scalds and bath water which is too hot is the biggest cause of fatal accidents to under fives.

PREVENTION:

Get a bath thermometer. Think about your home. Use oven guards, fireguards and electrical socket covers.

WHAT TO DO:

If your child has a minor burn run under cool water for 20 minutes. Do not apply butter or any fatty substance. Cover loosely with cling film or a clean plastic bag. Take your child to A&E if the burn is severe.

Eating a varied diet

Children should be encouraged to eat a varied diet. They should eat foods from each of the four main food groups every day. The four main food groups are:

- Bread, other cereals and potatoes.
- Fruit and vegetables.
- Milk and dairy foods.
- Meat, fish and alternatives such as pulses (peas, beans and lentils), eggs, vegetable proteins and soya.

A varied diet is associated with better health as it is more likely to contain all the vitamins and minerals the body needs.

What can I do?

Many parents are unaware of the dangers of childhood obesity but by following the top tips below you can make a difference to your child's health.

- 1. Sugar Swaps** - Swapping sugary snacks and drinks for ones that are lower in sugar can make a huge difference.
- 2. Meal Time** - It's important for kids to have regular, proper meals as growing bodies respond better to routine.
- 3. Snack Check** - Many snacks are full of the things that are bad for us - sugar, salt, fat and calories. So try and keep a careful eye on how many the kids are having.
- 4. Me Size Meals** - It's important to make sure kids get just the right amount for their age.
- 5. 5 A Day** - 5 portions of fruit and/or vegetables a day.
- 6. Cut Back Fat** - Too much fat is bad for us. It's not always easy to tell where it's lurking.
- 7. Up and About** - Most of us spend too long sitting down. Keep active. Encourage your child to walk, you may need to use child safety reins.

Source: www.nhs.uk/Start4Life
DoH 2009 (www.dh.gov.uk/obesity).



A healthy weight

A combination of the right food and exercise

Your Health Visitor will be able to discuss your child's weight with you to determine if they are under or overweight. If you are worried your child is underweight and they have a poor appetite, discuss with your Health Visitor or Doctor. If they are overweight and it is not treated early on it could cause health problems such as raised blood pressure and cholesterol, type 2 diabetes, early puberty, asthma and other respiratory illnesses.

Being overweight is rarely to do with a medical problem, many kids simply have unhealthy diets and don't do enough exercise. It is better to prevent your child becoming overweight (or obese) in the first place. A combination of good sleep patterns, a healthy varied diet and regular exercise will all keep your child to a healthy weight. It can be hard to recognise weight issues in our own children. It is important that parents spot the signs early on to prevent things getting worse and to promote healthy growth.

You are responsible for your child's health and well-being, this includes what they eat. A healthy varied diet and exercise is the simple answer to many worries about being overweight. Try to have family outings which include walking and cycling so you can all get fitter together. Being active burns more energy and the body then starts to use up its fat stores.

1

I am worried about my child's weight.

2

Do some exercise together as a family and find out more about healthy eating.

3

If you are worried discuss with your Doctor or your local Weight Management Service.

Dietician says

Salt and Sugar is added to nearly all processed products. Three-quarters of the salt and sugar we eat is already in the food, the rest is what we add to cooking or shake on our meals. Children need less than 5g of salt a day (2g sodium).



Juice drink
23g sugar
(5 teaspoons)



Cereal bars
8g sugar
(1.5 teaspoons)



Fromage frais
12.4g sugar
(2 teaspoons)



What can I do?

If someone you know is having difficulties, you could offer the following:

- A listening ear.
- Ideas to cope with problems.
- Encouragement to get help.
- Practical support (for example offering to babysit).

If the family is unable to cope and they do nothing to get help you need to report it. In an emergency call **999**.

Worried about a child?

Getting support

Very few adults harm children deliberately and most often, when harm does happen, families need support, not punishment or the removal of their children. Social Workers and other professionals get involved when parents may be unable to protect their child from harm and need some help. In some cases the Police Child Abuse Investigation Teams will work with Social Workers to help protect children and decide whether an offence has been committed.

If you believe a child is suffering or is likely to suffer harm, or if you are worried about your own behaviour and need support contact your local Children's Social Care Team (see page 53 for details). In an emergency call **999**.

When we suspect, witness or are told of a child that is being harmed, action should be taken to stop things getting worse. Even if you think an incident is just a one-off, your information could be very important. Long-term abuse can have a negative affect on a child for the rest of their lives.

1

There are many signs of abuse, from physical and emotional injury leading to changes in behaviour.

2

Make sure your child knows who they can share worries with if and when they need to.

3

Seek advice about what practical and emotional support is available from schools and Children's Centres.



Abuse at home

When we suspect abuse of a child in our own home, we can react in many different ways. We may feel guilt, anger, disbelief or denial. Some of these reactions can prevent getting help to a child who needs it. You may not tell others because you fear that the child will be at further risk of harm. You may love the person who is causing the harm and not want to believe what is happening. You need to put your child's safety first.



Unusual behaviour

Sometimes stress can lead to a 'tic' - a sudden, repetitive, non-rhythmic movement involving a distinct muscle group, like an uncontrolled eye blinking. Head banging or banging the head against a wall or cot on purpose is another common behaviour, especially in boys. They are not trying to be naughty or annoying. Your Doctor can advise if there are additional concerns. Once the Doctor gives your child a clean bill of health, the best thing to do is just ignore the tic, help minimise stress and make sure your child is getting enough sleep.

Good mental health

Support healthy social and emotional development

Responsive relationships help build positive attachments that support your baby's healthy social and emotional development. These relationships form the foundation of good mental health for infants, toddlers and pre-schoolers.

An infant's world is usually pretty small. Infants grow up among family, friends and community. They learn about themselves and the world around them through the relationships they have with the people in their lives. Babies who are made to feel loved and cherished learn that they are lovable. They then grow up feeling good about themselves and develop friendships more easily. You cannot spoil your baby by responding to their needs. By responding to their needs your baby will become more confident and independent in later years.

Stress is something as adults we come to accept and manage. Babies and young children are unable to recognise and cope with situations that cause stress in their lives. Instead, they can show their stress and anxiety in physical and emotional outbursts.

1

How do I bond with my baby?

2

Use eye contact, smile and hold them up so you are face to face.

3

This contact, support and security will help your baby develop in many ways.

What can I do?

- Surround your child with nurturing relationships.
- Be happy and show it.
- Create a trusting and loving environment.
- Provide stable and consistent caregivers at home and in childcare.
- Try to understand your child's cues and respond.
- Spend unhurried time together.
- Comfort and reassure.
- Respond to your baby.
- Learn about developmental stages and have appropriate expectations.
- Have good relationships and ways to manage conflict.
- Your actions and mood may affect your child.
- Identify early signs of emotional or mental problems.

Useful national contacts

Asthma UK

0800 121 62 44
www.asthma.org.uk

Baby LifeCheck

www.babylifecheck.co.uk

Child Accident Prevention Trust (CAPT)

020 7608 3828
www.capt.org.uk

ISIS

Advice on sleep
www.isisonline.org.uk

Meningitis Now

0808 80 10 388
www.meningitisnow.org

National Breastfeeding Helpline

0300 100 0212
www.breastfeedingnetwork.org.uk
email: enquiries@breastfeedingnetwork.org.uk

Netmums

Parenting advice and information in England, Wales, Scotland and Northern Ireland.
www.netmums.com

NHS Choices

www.nhs.uk



111

If you think you need help urgently during the day or night you should call 111 before you go to any other health service.

By calling 111 you will be directed straight away to the local service that can help you best. It is available 24 hours a day, 365 days a year and is free to call, including from a mobile. You should call NHS 111:

- When you need help fast but it's not life threatening.
- When you think you need to go to A&E or another NHS urgent care service.
- When it's outside of your Doctor's surgery hours.
- When you do not know who to call for medical help.
- If you do not have a local Doctor to call.

Call 999 in an emergency

Useful local contacts

More Life (for Islington children)

0113 812 5233
Email: team@more-life.co.uk

Islington Family Information Service

020 7527 5959 (weekdays 9am-4pm)
fis@islington.gov.uk
www.islington.gov.uk

Islington Breastfeeding and Weaning Mum-to-Mum Support

020 3316 3439

Children's Centres Islington

Ambler Children's Centre

Blackstock Road, N4 2DR.
020 7359 7628

Archway Children's Centre

Vorley Road, N19 5HE.
020 7527 4827

Bemerton Children's Centre

1a Coatbridge House, Carnoustie Drive
N1 0DX. 020 7527 4806

Conewood Street Children's Centre

14 Conewood Street, N5 1DL.
020 7527 4440

The Factory Children's Centre

107 Matthias Road, N16 8NP.
020 7923 9879

Golden Lane Children's Centre

Golden Lane Campus, Whitecross
Street, EC1Y 8JA. 020 7786 4800

Hungerford School and Children's Centre

Children's centre activities and services:
249 Hungerford Road, N7 9LF.
020 7619 9149

Childcare is based at our Goodinge
site: Corporation Street, N7 9EH.
020 7607 5850

We also have childcare and Children's
Centre services at: Robert Blair
Primary School and Nursery, Brewery
Road, N7 9QJ. 020 7697 4827

Kate Greenaway Nursery School and Children's Centre

York Way Court, Copenhagen Street,
N1 0UH. 020 7527 4850

Hornsey Road Children's Centre

8 Tiltman Place, Hornsey Road,
N7 7EN. 020 7527 2005

Margaret McMillan Nursery School and Children's Centre

Hornsey Rise, N19 3SF.
020 7281 2745

Moreland Children's Centre

Moreland Community House,
31 Moreland Street, EC1V 8BB.
020 7527 3500

New River Green Children's Centre

23 Ramsay Walk, N1 2SX.
020 7527 4813

North Islington Nursery School and Children's Centre

110-112 Tollington Park, N4 3RB.
020 7527 4844

Packington Children's Centre

Childcare services: 32a Popham Road,
N1 8SJ. 020 7226 6436
Family and community services:
190 Packington Square, N1 7UG.
020 7359 3302

Paradise Park Children's Centre

164 Mackenzie Road, N7 8SE,
020 7697 7330

Willow Children's Centre

1a Holbrooke Court, N7 0BF.
020 7527 1990



Breastfeeding Islington



@BFIslington



BF support Islington